arXiv:hep-th/0305214v1 24 May 2003

BELARUS NATIONAL ACADEMY OF SCIENCES
B.I. STEPANOV's INSTITUTE OF PHYSICS

Red'kov V.M.[1]

ON WKB-QUANTIZATION FOR KEPLER PROBLEM IN EUCLIDE, RIEMANN AND LOBACHEVSKY 3-SPACE

Quantum mechanical WKB-method is elaborated for the known quantum Kepler problem in curved 3-space models Euclide, Riemann and Lobachevsky in the framework of the complex variable function theory. Generalized Schrödinger, Klein-Fock hydrogen atoms are considered. Exact energy levels are found and their exactness is proved on the base of exploration into n-degree terms of the WKB-series. Dirac equation is solved too, but only approximate energy spectrum is established.

18 page, 68 references

---

[1] E-mail: redkov@dragon.bas-net.by

1. Introduction

It is well known that energy spectrum of the hydrogen atom had been calculated long before creating the consistent quantum mechanical theory. It was established that the Bohr-Sommerfeld rules, basis of the "old" quantum mechanics even without rigorous mathematical foundation, are closely referred to the so-called WKB-approximation it the consistent quantum theory [1-45].

Looking for solvable models in the framework of "new" quantum theory, some coolness towards approximate methods and any achievements of the Bohr-Sommerfeld mechanics was inevitable. But now and then the same question arises in the literature: why in the case of hydrogen atom the Bohr-Sommerfeld rule leads to the known exact energy spectrum. Also, from time to time in the literature one can face statement of the sort: in a potential φ the Bohr-Sommerfeld quantization gives an exact result $\epsilon_n(\varphi)$.

In the present work we turn again to a hydrogen atom but now placed concurrently in three different curved space backgrounds: Euclide E_3 (zero curvature), Riemann S_3 (positive constant curvature), and Lobachevsky H_3 (negative constant curvature). We have considered models based on Schrödinger, Klein-Fock equations in these three space models and shortly analyzed a hydrogen atom described by Dirac equation in E_3-space. From the very beginning it should be noted that these three hydrogen atom models prove themselves essentially different from mathematical and physical standpoint [46-68], and their study manifestly adds to general geometry-physics ground and to development of the quasi-classical method itself .

For the first time the hydrogen atom in a curved space-time was considered by E. Schrödinger [46]. In this work he elaborated the so-called factorization method as applied to some quantum mechanical eigenvalues problems, in particular the harmonic oscillator and the hydrogen atom. However, the method fails to function for a continuous part of the hydrogen energy spectrum. An idea to alter the task in order to obtain only discrete energy levels had arisen. Such a full discreteness is achieved by mere placing the hydrogen atom into a Riemannian 3-space of constant positive curvature S_3, because this 3-space is compact and therefore it might reproduce the effect of a finite dimensions box.

In hyper-spherical coordinates (χ, θ, ϕ) of S_3-model

$$dl^2 = d\chi^2 + \sin^2\chi\,(d\theta^2 + \sin^2\theta d\phi^2) \tag{1.1a}$$

the quantum mechanical Hamiltonian of the system has the form

$$H = -\frac{1}{2}\,\frac{1}{\sqrt{g}}\,\frac{\partial}{\partial x^\alpha}\,\sqrt{g}\,g^{\alpha\beta}\,\frac{\partial}{\partial x^\alpha}\;-\;\frac{e}{\tan\chi}\,. \tag{1.1b}$$

Let us use the following notation: the curvature radius ρ is a length unit, M is an electron mass, $\hbar^2/M\rho^2$ is an energy unit, $e = \frac{\alpha}{\rho}/\frac{\hbar^2}{M\rho^2}$ is an effective constant characterizing intensity of the Coulomb interaction. The sign at $(e/\tan\chi)$ hand-picked in eq. (1.1b) As expected the energy spectrum of the task turns out to be discrete in full, and as shown in [46] it is

$$\epsilon_n = -\frac{e^2}{2n^2} + \frac{1}{2}\,(n^2 - 1)\,, \quad n = 1, 2, 3, \ldots\,. \tag{1.1c}$$

The level ϵ_n displays n^2-degeneration similar to that existing in ordinary hydrogen atom theory. The number of negative energy levels is finite and it equals

$$x_0 - 1, \qquad \text{where} \qquad x_0^2 = \frac{1}{2}(1 + \sqrt{1 + 4e^2}), \tag{1.1d}$$

if x_0 is integer; at this there exists a zero level $\epsilon = 0$. When x_0 is fractional the number of negative energy levels equals an entire part of x_0. As long as $x_0 \geq 1$ holds, no matter how small is e (arbitrary small Coulomb attraction) always there will exists at least one negative energy level. In other words always there will exist at least one bound state of the system. It should be especially mentioned that in a E_3-based model of the hydrogen atom, no matter how small the intensity of the Coulomb attraction is there exists an infinite set of energies being concentrated near a zero point.

A hydrogen atom model in Lobachevsky space H_3 turns out to reveal still more uncommon properties [47]. In hyper-spherical coordinates of H_3-space

$$dl^2 = d\chi^2 + \sinh^2\chi(d\theta^2 + \sin^2\theta d\phi^2) \tag{1.2a}$$

the quantum mechanical Hamiltonian of the system looks

$$H = -\frac{1}{2}\,\frac{1}{\sqrt{g}}\,\frac{\partial}{\partial x^{\alpha}}\,\sqrt{g}\,g^{\alpha\beta}\,\frac{\partial}{\partial x^{\alpha}} \;-\; \frac{e}{\tanh\chi}\;. \tag{1.2b}$$

In [47] an energy spectrum and corresponding wave functions $\Psi_{\epsilon lm}(\chi,\theta,\phi)$ were found. The energy spectrum has a discrete and continuous parts; and the discrete part contains a finite number of energy levels

$$-\frac{e^2}{2} \le \epsilon \le (\frac{1}{2} - e) : \qquad \epsilon_n = -\frac{e^2}{2n^2} - \frac{1}{2}\,(n^2 - 1)\;, \quad n = 1,2,3,...,N\;. \tag{1.2c}$$

At this the level ϵ_n is n^2-degenerate in a way similar to the common hydrogen atom in the flat space. At $\epsilon \ge (\frac{1}{2} - e)$ the energy spectrum is continuous. The number N of discrete energy levels equals to $(\sqrt{e} - 1)$ at integer e, and N equals to an entire part of $\sqrt{e}$ if $\sqrt{e}$ is fractional. From the above it follows that along with decreasing e-parameter the number N gradually decreases too, and the point $e = 1$ represents a crucial moment after which no bound states in the system exists. In other words, in the H_3-space, the hydrogen atom is possible bound system only at $e > 1$.

Here one remark should be added. The distinctive features of the Kepler quantum problem in curved space backgrounds described above show potential value of these systems (and similar to them) from another view point. Any task, firstly being formulated as an old quantum problem in a new space background can be re-formulated as a certain special quantum problem in an old flat space background. So, there exists a systematic way based on geometry to reproduce some quantum mechanical potentials with physically interesting properties. Such a technique might be understood as a way to reproduce alternatively the usual potential approach to quantum mechanics through a geometry. Obviously, such an approach was discussed and used in the literature (for example, see [66]).

2. WKB-quantization and hydrogen atom in E_3-space

In Section 2, general mathematical structure of n-order term Q_n in WKB-series is established. It is proved that all terms $Q_n, n \ge 2$ do not contribute into an exact quantum mechanical quantization condition. In the light of this, success of the old quantum mechanics is fully rationalized.

Let us consider again a non-relativistic hydrogen atom in Euklide space model E_3. Schrödinger's equation after separation of variables by a substitution $\Psi = R(r)\,Y_{lm}(\theta,\phi)$ leads to

$$\frac{1}{r^2}\frac{d}{dr}r^2\frac{d}{dr}\,R \;+\; [\;\frac{2M}{\hbar^2}(\epsilon + \frac{\alpha}{r}) \;-\; \frac{l(l+1)}{r^2}\;]\,R \;= 0\;. \tag{2.1}$$

Let t be a new variable: $t = \ln r,\; r = e^t,\; t \in (-\infty, +\infty)$, then

$$\frac{d^2}{dt^2}R \;+\; \frac{d}{dt}R \;+\; [\;\frac{2\epsilon M\,e^{2t} + 2\alpha M\,e^{t}}{\hbar^2} \;-\; l(l+1)\;]R = 0\;. \tag{2.2a}$$

One can exclude the first derivative term by a substitution $R(t) = e^{-t/2}\,S(t)$

$$(\;\frac{d^2}{dt^2} \;+\; \frac{\Pi^2}{\hbar^2}\;)\,S = 0\;,$$

$$\Pi^2 = 2\epsilon M\,e^{2t} + 2\alpha M\,e^{t} - L^2\;, \qquad L^2 = \hbar^2(l + \frac{1}{2})^2 \tag{2.2b}$$

and further

$$S(t) = exp\,[\frac{i}{\hbar}\int Q(t)\,dt\,]\;, \qquad \frac{\hbar}{i}\frac{d}{dt}Q \;+\; Q^2 \;-\; \Pi^2 = 0\;. \tag{2.3}$$

Now let us expand the function $Q(t)$ into a series in terms of $(\hbar/i)^n$

$$Q(t) = \sum_{n=0}^{\infty}\,[\;(\frac{\hbar}{i})^n Q_n(t)\;]\;. \tag{2.4}$$

Substituting this formal decomposition into eq. (2.3b), we will have

$$(Q_0^2 - \Pi^2) \; + \sum_{n=1}^{\infty} \; (\frac{\hbar}{i})^n \; [\; \frac{d}{dt} \, Q_{n-1} \; + \sum_{k=0}^{n} Q_{n-k} \, Q_k \;] = 0 \; , \tag{2.5}$$

from which it follows

$$Q_0^2 = \Pi^2 \; , \;\; Q_n = - \, \frac{1}{2Q_0} \, (\, \frac{d}{dt} \, Q_{n-1} \; + \sum_{k=1}^{n-1} Q_{n-k} \, Q_k \;) = 0 \; , \quad n = 1, 2, ... \tag{2.6a}$$

Starting from the known Q_0, the relations (2.6a) enable us to calculate generally speaking all remaining Q_n; some first such terms are (below the symbol $'$ will denote (d/dt)-derivative):

$$Q_0^2 = \Pi^2 \qquad \Longrightarrow \qquad Q_0 = \pm \sqrt{\Pi^2} \; ,$$

$$Q_1 = - \, \frac{1}{2Q_0} \, Q_0' \; , \qquad Q_2 = - \, \frac{1}{2Q_0} \, (\, Q_1' \; + Q_1^2 \,) \; , \; ... \tag{2.6b}$$

Now we are at the point to formulate general quantization condition:

> We will assume that the wave function $S(t)$ corresponding to a bound state, being considered as a function of a complex variable t, has a finite numbers of zeros in the complex plane, which are allocated at real r-axis between classical turning points. According the known theorem in complex variable functions theory, the number of such zeros of $S(t)$ within certain domain can be calculated through an integral of the logarithmic derivative $(\ln S(t))'$ along a contour bounding that domain:

$$\frac{1}{2\pi i} \oint_{\mathcal{L}} [\; \frac{d}{dt} \; \ln S(t) \;] \; dt = n \; , \tag{2.7}$$

Taking into account (2.3), the quantization condition (2.7) will take the form

$$\oint_{\mathcal{L}} \; Q(t) \; dt = 2\pi\hbar \; n \; , \tag{2.8}$$

from which substituting a series (2.4) instead of $Q(t)$, we arrive at

$$\sum_{n=0}^{\infty} [\; (\frac{\hbar}{i})^n \; \oint_{\mathcal{L}} \; Q_n(t) \; dt \;] \; = 2\pi\hbar \; n \; . \tag{2.9a}$$

It should be especially emphasized that relations (2.7), (2.8) and (2.9a) are precise mathematical conditions without any approximation.

Below, at calculating separate contour integrals in (2.9a) we will be needing to use a complex variable $z = e^t$, so instead of (2.9a) one has

$$\sum_{n=0}^{\infty} [\; (\frac{\hbar}{i})^n \; \oint_{\mathcal{L}} \; Q_n(z) \; \frac{dz}{z} \;] \; = 2\pi\hbar \; n \; . \tag{2.9b}$$

All terms $z^{-1} \, Q_n(z)$ in the WKB-series (2.9b) are single valued and analytical functions of the variable z just in 3-connected domain of z-plane, bounded by three contours $\mathcal{L}^{all} = \mathcal{L}_0 + \mathcal{L}_\infty + \mathcal{L}$ (see. Fig. 1; dashed line designates a special cut needed to allow for existence of two branch points of $Q_0 = \sqrt{\Pi^2}$; $\mathcal{L}_0$ is any enough small contour around zero point $z = 0$; $\mathcal{L}_\infty$ is any enough large contour around ∞-point):



Fig. 1 (3-connected domain and Cauchy theorem)

Besides, all functions $z^{-1}\,Q_n(z)$ are continuous up to the boundary $\mathcal{L}^{all}$, hense Cauchy theorem will hold in this 3-connected domain:

$$- \oint_{\mathcal{L}} Q_n(z)\,\frac{dz}{z} \;+\; \oint_{\mathcal{L}(0)} Q_n(z)\,\frac{dz}{z} \;+\; \oint_{\mathcal{L}(\infty)} Q_n(z)\,\frac{dz}{z} \;=0\;; \qquad (2.10a)$$

where the sign minus before the first integral refers to "improper" direction of going around the contour $\mathcal{L}$. With the use of conventional residue notation, from (2.10a) we arrive at the main working formula

$$\oint_{\mathcal{L}} Q_n(z)\,\frac{dz}{z} \;=-\;(2\pi i)\sum_{z=0,\infty} res\;\frac{Q_n(z)}{z}\;. \qquad (2.10b)$$

Thus, for the zero order term in (2.9b) we obtain

$$\oint_{\mathcal{L}} Q_0(z)\,\frac{dz}{z} = -\;(2\pi i)\sum_{z=0,\infty} res\;\frac{\sqrt{2\epsilon M\,z^2+2\alpha M\,z-L^2}}{z} =$$

$$= 2\pi\;(\;-L+i\;\frac{\alpha M}{\sqrt{2\epsilon M}}\;)\;. \qquad (2.11)$$

As for the next order term $Q_1(t)$ we have

$$Q_1(t) = -\;\frac{1}{2Q_0(t)}\;Q_0'(t) = -\frac{1}{2}\;\frac{(2\epsilon M\;e^{2t}+\alpha M\;e^t)}{(2\epsilon M\;e^{2t}+2\alpha M\;e^t-L^2)}\;, \qquad (2.12a)$$

and its contribution is

$$\frac{\hbar}{i}\oint_{\mathcal{L}} Q_1(z)\;\frac{dz}{z} = -\;\pi\,\hbar\;. \qquad (2.12b)$$

With regard to Q_0 and Q_1, eq. (2.9b) gives the following ($L=\hbar\;(l+\frac{1}{2})$)

$$-L+i\;\frac{\alpha M}{\sqrt{2\epsilon M}}-\frac{\hbar}{2}=\hbar\;n \qquad\Longrightarrow\qquad \epsilon=-\frac{\alpha^2 M}{2\hbar^2 N^2}\;. \qquad (2.13)$$

where $N=(n+l+1)$. The energy spectrum obtained coincides with the known exact result from the Schrödinger quantum mechanical equation.

Now it is the point to understand this coincidence from a more general viewpoint. What are the reasons for such a success of the seemingly approximate technics. The form of Q_0 as a second order polynomial is relevant point; and another essential fact is that $\mathcal{L}$-contour encloses zeros (classical turning points or branch points of $\sqrt{P}$) of this polynomial. At these general assumptions we always will obtain

$$Q_0(t)=\sqrt{P}\;,\qquad P=(Ae^{2t}+Be^t+C)\;, \qquad (2.14)$$

$$res\;_{z=0}\frac{Q_0(z)}{z}=\sqrt{C}\;,\qquad res\;_{z=\infty}\frac{Q_0(z)}{z}=-\;\frac{B}{2\sqrt{A}}\;\Longrightarrow$$

$$\oint_{\mathcal{L}} Q_0\;dt=2\pi\;[-\sqrt{-C}\;+i\;\frac{B}{2\sqrt{A}}\;]\;;$$

$$Q_1(t)=-\frac{Q_0'}{2Q_0}=-\frac{1}{2}\;\frac{P'}{2\sqrt{P}}\;\frac{1}{\sqrt{P}}=-\frac{1}{2}\;\frac{P'}{2P}\;,\qquad\Longrightarrow$$

$$\frac{\hbar}{i}\oint_{\mathcal{L}} Q_1\;dt=-\frac{\hbar}{2}\;2\pi\;.$$

Now we are going to elucidate general structure of all terms of greater order than the first. For $Q_2(t)$ we have

$$Q_2(t)=-\frac{1}{2Q_0}\;(\;Q_1'+Q_1^2\;)=$$

$$=-\;\frac{1}{2\sqrt{P}}\;[(\;(-\frac{1}{2})\frac{P''}{2P}+(\frac{1}{2})\;\frac{P'P'}{2P^2})+(\frac{1}{4})\;\frac{P'P'}{2^2P^2}\;]=$$

$$= - \frac{1}{2\sqrt{P}} [- \frac{P''}{2^2 P} + \frac{5}{4} \frac{P'P'}{2^2 P^2}] \, . \tag{2.15}$$

Further it will be helpful to use special designation. The terms Q_1, Q_2 let be expressed symbolically as

$$Q_1 \sim \frac{2'}{2} \, , \qquad Q_2 \sim \frac{1}{\sqrt{P}} (\frac{2'}{2} \oplus \frac{4'}{4}) \, , \tag{2.16a}$$

where

$$(-\frac{P'}{2P}) \text{ refers to } \frac{2'}{2} \, ,$$
$$(- \frac{P''}{2^2 P}) \text{ refers to } \frac{2'}{2} \, ,$$
$$(\frac{1}{4}) \frac{P'P'}{2^2 P^2} \text{ refers to } \frac{4'}{4} \, .$$

In other words, the notation used keeps only information relevant to further calculating the residues res_0 and res_∞:

1) we indicate the power of polynomials at numerator and denominator of any term;

2) also we indicate that the numerator - polynomial in e^t was differentiated with respect to t-variable and therefore the constant term of the polynomial is suppressed.

No matter how many times the polynomial was differentiated, the same single symbol $'$ stands for all these cases.

In such symbolic notation the operator d/dt acts as follows

$$\frac{d}{dt} \frac{1}{\sqrt{P}} = - \frac{1}{2\sqrt{P}} \frac{P'}{P} \sim \frac{1}{\sqrt{P}} \frac{2'}{2} \, ,$$

$$\frac{d}{dt} \frac{2'}{2} \sim [\frac{2'}{2} \oplus \frac{4'}{4}] \, ,$$

$$\frac{d}{dt} \frac{4'}{4} \sim [\frac{4'}{4} \oplus \frac{6'}{6}] \, , \ldots$$

Now we are ready to proceed to Q_3 term:

$$Q_3 = - \frac{1}{2Q_0} (Q_2' + 2Q_1 Q_2) \sim$$

$$\sim \frac{1}{\sqrt{P}} \left\{ [\frac{d}{dt} \frac{1}{\sqrt{P}} (\frac{2'}{2} \oplus \frac{4'}{4})] + \frac{2'}{2} \frac{1}{\sqrt{P}} (\frac{2'}{2} \oplus \frac{4'}{4}) \right\} \sim$$

$$\sim \frac{1}{P} \left[\frac{2'}{2} (\frac{2'}{2} \oplus \frac{4'}{4}) \oplus \frac{d}{dt} (\frac{2'}{2} \oplus \frac{4'}{4}) \oplus \frac{2'}{2} (\frac{2'}{2} \oplus \frac{4'}{4}) \right] \, .$$

So, the final structure of Q_3 looks as

$$Q_3 \sim \frac{1}{P} (\frac{2'}{2} \oplus \frac{4'}{4} \oplus \frac{6'}{6}) \, . \tag{2.16b}$$

Let us consider else one term:

$$Q_4 = - \frac{1}{2Q_0} (Q_3' + 2Q_1 Q_3 + Q_2^2) \, .$$

With the use of symbolic notation we get to

$$Q_4 \sim \frac{1}{\sqrt{P}} \{ \frac{d}{dt} [\frac{1}{P} (\frac{2'}{2} \oplus \frac{4'}{4} \oplus \frac{6'}{6})] \oplus \qquad \text{(2.16c))}$$

$$\oplus \frac{2'}{2} \frac{1}{P} (\frac{2'}{2} \oplus \frac{4'}{4} \oplus \frac{6'}{6}) \oplus [\frac{1}{\sqrt{P}} (\frac{2'}{2} \oplus \frac{4'}{4})]^2 \} \sim$$

$$\sim \frac{1}{P\sqrt{P}} \, [\, \frac{2'}{2} \oplus \frac{4'}{4} \oplus \frac{6'}{6} \oplus \frac{8'}{8} \,] \, .$$

At this point we are ready to write down symbolically the following general structure of an arbitrary order term Q_n of WKB-series:

$$Q_n = \frac{1}{(\sqrt{P})^{n-1}} \left[\frac{2'}{2} \oplus \frac{4'}{4} \oplus ... \oplus \frac{(2n)'}{(2n)} \right] . \tag{2.17a}$$

It is readily verified that all required residues vanish identically:

$$res_{\;\; z=0} \frac{Q_n(z)}{z} = \frac{1}{(\sqrt{C})^{n-1}} \, 0 = 0 \;\; ,$$

$$res_{\;\; z=\infty} \frac{Q_n(z)}{z} = -res_{\;\; y=0} \frac{Q_n(y^{-1})}{y} = \tag{2.17b}$$

$$= - \, res_{\;\; y=0} \frac{1}{y \; (\; \sqrt{Ay^{-2} + By^{-1} + C} \;)^{n-1}} [\frac{2'}{2} \oplus \frac{4'}{4} \oplus ... \oplus \frac{(2n)'}{(2n)}]_{z=y^{-1}} =$$

$$= - \, res_{\;\; y=0} \frac{y^{n-1}}{y \; (\; \sqrt{A + By^{-1} + Cy^{-2}} \;)^{n-1}} \;\; const \;\; = 0 \, . \qquad (2.17c))$$

So, the success of Borh-Sommerfeld quantization for the hydrogen atom is rationalized in full.

3. Non-relativistic hydrogen atom in Lobachevsky space H_3

In Section 3 it is shown that the Borh-Sommerfeld quantization as applied to the hydrogen atom in Lobachevsky space H_3 provides us with an exact energy spectrum of this generalized atom model.

In Lobachevsky space the radial wave equation has the form

$$\frac{1}{\sinh^2} \frac{d}{d\chi} \sinh^2 \chi \frac{d}{d\chi} \, R + [\, \frac{2M\rho^2}{\hbar^2} \, (\epsilon + \frac{\alpha}{\rho \tanh \chi}) - \frac{l(l+1)}{\sinh^2 \chi} \,] \, R(\chi) = 0 \, . \tag{3.1}$$

Introducing a new variable $t = \ln(\tanh \chi)$

$$\frac{d^2}{dt^2} R + \frac{d}{dt} R + \frac{1}{(1-e^{2t})^2} [\frac{2\epsilon\rho^2 M e^{2t} + 2\alpha\rho^2 M e^t}{\hbar^2} - l(l+1)(1-e^{2t})] R = 0 \tag{3.2a}$$

and excluding the first-derivative term by the substitution $S = e^{t/2} \, R$ one gets to

$$\frac{d^2}{dt^2} \, S \; + \; \left[\, \frac{1}{(1-e^{2t})^2} \, \left(\frac{2\epsilon\rho^2 M \; e^{2t} + 2\alpha\rho^2 M \; e^t}{\hbar^2} \; - \; l(l+1)(1-e^{2t}) \right) - \frac{1}{4} \, \right] \; S = 0 \, . \tag{3.2b}$$

Further we are forced to move through trial and error method. Because we have known an exact energy spectrum of the system in advance, we are going to proceed in the following way: in the exact quantization condition let us take into account only two first terms of a certain formal WKB-series, on obtaining the required exact result we will analyze next terms.

Firstly, with the use of the trick

$$- \, \frac{l(l+1)}{1-e^{2t}} = \frac{1}{1-e^{2t}} [\, -(l+\frac{1}{2})^2 + \frac{1}{4} \,] = \frac{1}{1-e^{2t}} \, (-\frac{L^2}{\hbar^2} + \frac{1}{4}) \; ;$$

eq. (3.2b) becomes

$$\frac{d^2}{dt^2} S + \left[\, \frac{2\epsilon\rho^2 M \; e^{2t} + 2\alpha\rho^2 M \; e^t - L^2(1-e^{2t})}{\hbar^2 (1-e^{2t})^2} \; + \frac{1}{4}(\frac{1}{1-e^{2t}} - 1) \right] \; S = 0 \, , \tag{3.3a}$$

or shorter

$$\frac{d^2}{dt^2} \, S(t) \; + \; [\, \frac{\Pi^2(t)}{\hbar^2} \; + \; \Delta(t) \,] \, S(t) = 0 \, , \tag{3.3b}$$

$$\Pi^2(t) = \frac{(2\epsilon\rho^2 M + L^2)e^{2t} + 2\alpha\rho^2 M\ e^t - L^2}{(1-e^{2t})^2}\ , \qquad \Delta(t) = \frac{1}{4}\frac{e^{2t}}{(1-e^{2t})}\ .$$

Further we act as in Section 2:

$$S(t) = exp[\ \frac{i}{\hbar} \int Q(t)dt\ ]\ , \qquad \frac{\hbar}{i}\ \frac{d}{dt}Q + Q^2 - \Pi^2 + (\frac{\hbar}{i})^2\Delta = 0\ ,$$

$$Q(t) = \sum_{n=0}^{\infty} [\ (\frac{\hbar}{i})^n\ Q_n(t)\ ]\ , \qquad Q_0 = \sqrt{\Pi^2}\ ,\ Q_1 = -\frac{1}{2Q_0}\ Q_0'\ ,$$

$$Q_2 = -\frac{1}{2Q_0}\ [\ Q_1' + Q_1^2 + \Delta\ ]\ ,...$$

$$Q_n = -\ \frac{1}{2Q_0}\ [\ \frac{d}{dt}Q_{n-1}\ +\ \sum_{k=1}^{n-1} Q_{n-k}Q_k\ ] = 0\ ,\ n = 3,4,5,... \tag{3.4a}$$

Quantum condition is the same

$$\frac{1}{2\pi i}\ \oint_{\mathcal{L}} [\frac{d}{dt}\ln S(t)]\ dt = n \quad \Longrightarrow$$

$$\sum_{n=0}^{\infty} [\ (\frac{\hbar}{i})^n\ \oint_{\mathcal{L}} Q_n(t)\ dt\ ]\ = 2\pi\hbar\ n\ . \tag{3.4b}$$

In contrast to previous case (Section 2) here we are to employ Cauchy theorem for a 5-connected domain:



Fig. 2 (5-connected domain and Cauchy theorem)

Cauchy theorem provides us with relation

$$-\ \oint_{\mathcal{L}} Q_n(z)\ \frac{dz}{z} + \tag{3.5a}$$

$$+\ \oint_{\mathcal{L}(0)} Q_n(z)\ \frac{dz}{z}\ +\ \oint_{\mathcal{L}(-1)} Q_n(z)\ \frac{dz}{z} +$$

$$+\ \oint_{\mathcal{L}(+1)} Q_n(z)\ \frac{dz}{z}\ +\ \oint_{\mathcal{L}(\infty)} Q_n(z)\ \frac{dz}{z}\ = 0\ ;$$

the contour $\mathcal{L}$ is going around in improper direction which reflects the sing minus at the first term. From (3.5a) it follows

$$\oint_{\mathcal{L}} \frac{Q_n(z)}{z}\ dz\ = (-2\pi i)\ \sum res\ _{z=0,\pm1,\infty}\ (\frac{Q_n(z)}{z})\ . \tag{3.5b}$$

Two first terms of the WKB-series contribute to quantum condition (to do formulae shorter dimensionless quantities are used)

$$\oint_{\mathcal{L}} Q_0(t)\ dt = (-2\pi i)\ \left[\ \sqrt{-L^2}\ -\ \frac{1}{2}\ (\ \sqrt{2\epsilon + 2e} + \sqrt{2\epsilon - 2e}\ )\right]\ ,$$

$$\frac{\hbar}{i}\oint_{\mathcal{L}} Q_1\frac{dz}{z} = -\pi\,\hbar\ . \tag{3.6}$$

which results in

$$\frac{i}{2}\,(\sqrt{2\epsilon+2e}+\sqrt{2\epsilon-2e}) = \hbar\,(n+l+1) \qquad \Longrightarrow$$

$$N = n+l+1, \qquad \epsilon = -\,\frac{e^2}{2N^2}-\frac{N^2}{2}\ . \tag{3.7a}$$

However we have known an exact energy formula [47]

$$\epsilon = -\,\frac{e^2}{2N^2}-\frac{N^2-1}{2}\ . \tag{3.7b}$$

So, (3.7a), what we have found, slightly differs from expected eq. (3.7b); difference is in an additive term $\frac{1}{2}$. Returning to the expression (3.3b) for Π^2, one notices that it would suffice a single change $(2\epsilon+L^2) \implies (2\epsilon-\hbar^2+L^2)$, to reproduce through 2-term WKB-quantization just the exact energy formula (3.7b). This change would lead to

$$\Delta = \frac{e^{2t}}{4(1-e^{2t})} \qquad \Longrightarrow \qquad (\frac{e^{2t}}{4(1-e^{2t})}+\frac{e^{2t}}{(1-e^{2t})^2}) = \frac{5-e^{2t}}{4}\ \frac{e^{2t}}{(1-e^{2t})^2}\ .$$

In other words, instead formulas (3.3b) one should begin with the following equations

$$\frac{d^2}{dt^2}\,S(t)\ +\ [\ \frac{\Pi^2(t)}{\hbar^2}\ +\ \Delta(t)\ ]\ S(t) = 0\ , \tag{3.8a}$$

$$\Pi^2(t) = \frac{(2\epsilon-\hbar^2+L^2)e^{2t}+2\alpha\,e^t-L^2}{(1-e^{2t})^2}\ ,$$

$$\Delta(t) = \frac{5-e^{2t}}{4}\ \frac{e^{2t}}{(1-e^{2t})^2}\ . \tag{3.8b}$$

For the following it will be convenient to have solved the task (3.8b) in a general form

$$Q_0(t) = \sqrt{\Pi^2(t)} = \frac{\sqrt{Ae^{2t}+Be^t+C}}{1-e^{2t}} = \frac{\sqrt{P(t)}}{1-e^{2t}}\ ,$$

and further

$$res\ _{z=0}\ \frac{Q_0}{z} = \sqrt{C}\ , \qquad res\ _{z=\infty}\ \frac{Q_0}{z} = 0\ ,$$

$$res\ _{z=+1}\ \frac{Q_0}{z} = -\ \frac{1}{2}\ \sqrt{A+B+C}\ ,$$

$$res\ _{z=-1}\ \frac{Q_0}{z} = -\ \frac{1}{2}\ \sqrt{A-B+C}\ ,$$

$$\oint Q_0\frac{dz}{z} = 2\pi[\ -\sqrt{-C}+\frac{i}{2}\ (\sqrt{A+B+C}+\sqrt{A-B+C})\ ]\ , \tag{3.9a}$$

$$Q_1 = -\ \frac{1}{2}(\frac{P'}{2P}+\frac{2e^{2t}}{1-e^{2t}})\ , \qquad \frac{\hbar}{i}\oint Q_1\frac{dz}{z} = -\ 2\pi\ \frac{\hbar}{2}\ . \tag{3.9b}$$

The 2-term WKB-quantization provides us with

$$\Big[\ -\sqrt{-C}+\frac{i}{2}\ (\sqrt{A+B+C}+\sqrt{A-B+C})\ \Big]\ -\ \frac{\hbar}{2} = \hbar\ n\ , \tag{3.9c)}$$

as applied to (3.8b) this leads to

$$i\ \frac{1}{2}\ \left(\sqrt{2\epsilon-1+2e}+\sqrt{2\epsilon-1-2e}\right) = N\ , \ \ N = (n+l+1)\ ,$$

from which it follows an exact energy formula. It can be readily verified by a straightforward calculation that the next term Q_2 (see (3.4a) and (3.8b)) gives a zero to the main quantization condition[2]

---

[2] General task of proving that all terms greater order than Q_2 do not contribute to the quantum condition looks quite interesting from mathematical viewpoint, but it will be omitted here.

4. Hydrogen atom in Riemann space of constant curvature S_3

To have analyzed the hydrogen atom in the same line on the Riemann constant curvature background we need no new ideas in addition to previous case (Section 3). And so it suffices to write down only several main relations:

Radial wave equation has the form

$$\frac{1}{\sin^2}\,\frac{d}{d\chi}\sin^2\chi\frac{d}{d\chi}\;R+[\frac{2M\rho^2}{\hbar^2}\;(\epsilon+\frac{\alpha}{\rho\tan\chi})-\frac{l(l+1)}{\sin^2\chi}\;]\;R(\chi)=0\;.$$

Quantization condition remains the same. Computation of contour integrals is reduced to four residues:

$$\oint_{\mathcal{L}}\frac{Q_n(z)}{z}\;dz\;=(-2\pi i)\sum\;res_{\;z=0,\pm i,\infty}\;(\frac{Q_n(z)}{z})\;.$$

2-term quantization gives

$$\frac{i}{2}\;(\sqrt{-(2\epsilon\rho^2M+\hbar^2)+2i\alpha\rho M}+\sqrt{-(2\epsilon\rho^2M-2i\alpha\rho M})=\hbar\;(n+l+1)\;,$$

from which it follows the exact energy formula

$$\epsilon=-\;\frac{e^2}{2N^2}+\frac{N^2-1}{2}\;.$$

5. WKB-analysis of the Klein-Fock equations in spaces E_3, H_3, S_3

The hydrogen atom models based on Klein-Fock equation in three spaces E_3, H_3, S_3 are described by radial equations which come within the general types that have been considered at non-relativistic case (see (2.17) and (3.9)), hence we can act nearly formally without any serious additional work. At this line one can readily produce three known exact energy spectra on the base of the 2-term quantization.

In flat space E_3

$$\epsilon=mc^2\left(1+\frac{\alpha^2/\hbar^2c^2}{N^2}\right)^{-1/2}\;. \tag{5.1}$$

In the Riemann and Lobachevsky spaces (upper and lower sign correspondingly)

$$\epsilon=mc^2\left(1+\frac{\alpha^2/\hbar^2c^2}{N^2}\right)^{-1/2}\;\sqrt{1\pm\frac{\hbar^2}{m^2c^2\rho^2}(N^2-1+\alpha^2/\hbar^2c^2)} \tag{5.3}$$

where $N=(n_r+1/2+\sqrt{(l+1/2)^2-\alpha^2/\hbar^2c^2}\;.$

6. Energy spectrum of a free Dirac particle on the sphere S_3

Omitting all details on separating the variables in the free Dirac equation in space of positive constant curvature S_3, let us start with a corresponding radial equation

$$\frac{d^2}{d\chi^2}f\;+\;[\;\frac{\rho^2(\epsilon^2-m^2c^4)}{c^2\hbar^2}-\frac{k^2+k\cos\chi}{\sin^2\chi}\;]\;f=0 \tag{6.1}$$

With a new variable $t=\ln\;\tan\chi$, eq. (6.1) takes on the form

$$\frac{d^2}{dt^2}\;f\;-\;\frac{1-e^{2t}}{1+e^{2t}}\;\frac{d}{dt}\;f\;+$$

$$+\frac{1}{\hbar^2}\;\frac{[\frac{\rho^2(\epsilon^2-m^2c^4)}{c^2}-L^2]e^{2t}-L^2}{(1+e^{2t})^2}\;f-\frac{k}{(1+e^{2t})^{3/2}}\;f=0\;. \tag{6.2}$$

Further let us act in accordance with a standard scheme:

$$f = exp[\frac{i}{\hbar}\int Q(t)dt] \; , \qquad Q(t) = \sum_{k=0}^{k=\infty} (\frac{\hbar}{i})^k Q_k(t) \; ,$$

$$\frac{1}{2\pi i} \oint_{\mathcal{L}} \frac{V'}{V} dt = n \; , \qquad \sum_{k=0}^{k=\infty} (\frac{\hbar}{i})^k \oint_{\mathcal{L}} Q_k(t) dt = 2\pi\hbar n$$

For two first terms in the quasi-classical series we have expressions:

$$Q_0(t) = \frac{\sqrt{[\frac{\rho^2}{c^2}(\epsilon^2 - m^2c^4) - L^2]e^{2t} - L^2}}{1 + e^{2t}}$$

$$Q_1(t) = -\frac{1}{2}\left(\frac{Q_0'}{Q_0} - \frac{1 - e^{2t}}{1 + e^{2t}}\right) = -\frac{1}{2}\left[\frac{[\frac{\rho^2}{c^2}\,(\epsilon^2 - m^2c^4) - L^2]\; e^{2t}}{[\frac{\rho^2}{c^2}\,(\epsilon^2 - m^2c^4) - L^2]e^{2t} - L^2} - 1\right] \; ,$$

their contributions are

$$\oint_{\mathcal{L}} Q_0 dt = 2\pi[-k\hbar + \sqrt{\frac{\rho^2}{c^2}(\epsilon^2 - m^2c^4)}] \; ,$$

$$\oint_{\mathcal{L}} Q_1 dt = 2\pi(-\frac{\hbar}{2}) \tag{6.3}$$

So, 2-term quantization gives

$$\sqrt{\frac{\rho^2}{c^2}(\epsilon^2 - m^2c^4)} = (n + k + \frac{1}{2}) \quad \Longrightarrow$$

$$\epsilon = \sqrt{m^2c^4 + \frac{c^2\hbar^2}{\rho^2}(n + k + \frac{1}{2})^2} \tag{6.4}$$

which is an exact energy spectrum for a free Dirac particle on the sphere S_3.

7. Dirac equation in spaces H_3 and S_3

Let us consider quasi-classical description of the hydrogen atom on the base of Dirac equation in H_3-space. After separation of variables in hyper-spherical coordinates the radial system is

$$\frac{df}{d\chi} + \frac{k}{\sinh\chi} f - \frac{\rho}{\hbar c}(\epsilon + mc^2 + \frac{\alpha}{\rho}\tanh^{-1}\chi) g = 0 \; ,$$

$$\frac{dg}{d\chi} - \frac{k}{\sinh\chi} g + \frac{\rho}{\hbar c}(\epsilon - mc^2 + \frac{\alpha}{\rho}\tanh^{-1}\chi) f = 0 \; , \tag{7.1}$$

so the task is reduced to a second order differential equation

$$\frac{d^2}{d\chi^2}\, f \; + \; \frac{(\alpha/\rho)}{\sinh\chi}\, \frac{1}{[\frac{\alpha}{\rho}\cosh\chi + (\epsilon + mc^2)\sinh\chi]}\; \frac{d}{d\chi}\, f \; + \tag{7.2}$$

$$+ \frac{\rho^2}{\hbar^2 c^2}\left[(\epsilon + \frac{\alpha}{\rho}\tanh^{-1}\chi)^2 \; - m^2c^4 \right] - \frac{k^2 + k\cosh\chi}{\sinh^2\chi}\, f \; +$$

$$+ \frac{k}{\sinh^2\chi}\; \frac{(\alpha/\rho)}{\frac{\alpha}{\rho}\cosh\chi + (\epsilon + mc^2)\sinh\chi}\, f = 0 \; .$$

With a new variable $\ln(\tanh(\chi/2)) = t$ this equation takes on the form

$$\frac{d^2}{dt^2} f \; + \; \left[\frac{1 - e^{2t}}{e^{2t} + 2ae^t + 1} - \frac{1 + e^{2t}}{1 - e^{2t}}\right] \frac{d}{dt} f \; +$$

$$+ \frac{1}{\hbar^2}\, \frac{1}{(1 - e^{2t})^2}\; \left[(2e^t\, \frac{\epsilon\rho}{c} \; + \; \frac{\alpha}{c}\,(1 + e^{2t}))^2 \; - \; m^2\rho^2c^4 4e^{2t} \; - \; L^2\,(1 - e^{2t})^2\right] \; -$$

$$- \, k \; \frac{2e^{t}}{1-e^{2t}} \; \frac{a(1+e^{2t})+2e^{t}}{(1+e^{2t})+a2e^{t}} \; f = 0 \; ; \tag{7.3a}$$

where

$$L^{2} = \hbar^{2} k^{2} \; , \qquad a = \frac{\epsilon + mc^{2}}{\alpha/\rho} \tag{7.3b}$$

or in a short form

$$f'' + V \; f' + (\frac{\Pi^{2}}{\hbar^{2}} + \Delta) f = 0 \; . \tag{7.4a}$$

Further acting upon old scheme:

$$f = exp[\frac{i}{\hbar} \int Q(t) dt] \; ,$$

$$\frac{\hbar}{i} Q' + Q^{2} + \frac{\hbar}{i} V \; Q - \Pi^{2} + (\frac{\hbar}{i})^{2} \Delta = 0 \tag{7.4b}$$

for terms of quasi-classical series we find recursive relations:

$$Q(t) = \sum_{n=0}^{n=\infty} (\frac{\hbar}{i})^{n} Q_{n}(t) \; \; :$$

$$Q_{0}^{2} = \Pi^{2} \; , \; Q_{1} = - \frac{1}{2Q_{0}} (Q_{0}' + V \; Q_{0}) \; , \; \; ... \tag{7.4c)}$$

Quantum condition is

$$\frac{1}{2\pi i} \oint_{\mathcal{L}} \frac{f'}{f} dt = n \quad \Longrightarrow \quad \sum_{k=0}^{k=\infty} (\frac{\hbar}{i})^{k} \oint_{\mathcal{L}} Q_{k}(t) dt = 2\pi\hbar \; N \tag{7.4d}$$

Here we need to discuss some new peculiarities. The matter is that the term Q_0 in $z = \tanh \frac{\chi}{2}$ variable has the form

$$Q_{0}(z) = \Pi(z) = \frac{1}{1-z^{2}} \; \sqrt{[2z \; \frac{\epsilon\rho}{c} + \frac{\alpha}{c}(1+z^{2})]^{2} - m^{2}\rho^{2}c^{2} \; 4z^{2} - L^{2}(1-z^{2})^{2}} \; , \tag{7.5}$$

which means that under square root stays 4-order polynomial. To make use the conventional residue theory we are forced to take for integration only such contours that enclose all four branch points of $Q_0(z)$-term.

Let us show that two zeros of the function $Q_0(z)$ are allocated in physical region and two zeros are allocated in non-physical one. To understand this let us consider in some detail such a relation

$$\tanh \chi = \frac{2 \tanh \frac{\chi}{2}}{1 + \tanh^{2} \frac{\chi}{2}} \qquad \Longrightarrow \qquad w^{2} + 1 - \frac{2w}{\tanh \chi} = 0 \; , \qquad w = \tanh \frac{\chi}{2};$$

which has two solutions:

$$w_{1} = z = \frac{\cosh \chi - 1}{\sinh \chi} \; , \qquad w_{2} = Z = \frac{\cosh \chi + 1}{\sinh \chi} \; .$$

Taking into account relations

$$z = 1 \; - 2 \; \frac{1 - e^{-\chi}}{e^{+\chi} - e^{-\chi}} \; < \; 1 \; , \; z = \tanh \frac{\chi}{2} \; ,$$

$$Z = 1 \; + 2 \; \frac{1 + e^{-\chi}}{e^{+\chi} - e^{-\chi}} \; > \; 1 \; , \qquad Z \neq \tanh \frac{\chi}{2} \; ;$$

one concludes that only one solution, z can be related with a physical variable $\tanh \frac{\chi}{2}, \chi \in (0, +\infty)$; whereas the second solution $Z \in (+1, +\infty)$ cannot be related in any sense to physical variable χ. Besides, between roots z and Z there exists simple connection:

$$w_{1} \; w_{2} = z \; Z = 1 \; . \tag{7.6}$$

Now let us compute what the term Q_0 gives into quantum condition. (7.4d):

$$\oint_{\mathcal{L}} Q_0(t)dt \ ,$$

integration should be taken along the contour enclosing classical turning points on positive χ-axis as shown in Fig. 3

Fig 3. Physical integration contour $\mathcal{L}(z)$.



However on mathematical reasons it will be convenient to change this integration contour $\mathcal{L}$ into doubled one $\mathcal{L}'$ (enclosing non-physical region of χ-variable)

$$\oint_{\mathcal{L}} Q_0(t)dt = \frac{1}{2} \oint_{\mathcal{L}'} Q_0(t)dt \ ; \qquad (7.6c))$$

Fig. 4. Doubled integration contour $\mathcal{L}'(z)$



With such a trick we will count up total number of zeros: physical and unphysical ones.
In so doing we will find contribution of the term Q_0 through conventional residue-technics:

$$\frac{1}{2} \oint_{\mathcal{L}'} Q_0(t)dt = (\frac{1}{2}) \ (-2\pi i) \sum_{0,\infty,\pm 1} res \ \frac{Q_0(z)}{z} \ . \qquad (7.7a)$$

Allowing for

$$res \ _{z=0} \ \frac{Q_0(z)}{z} = Q_0(z=0) = \sqrt{\frac{\alpha^2}{c^2} - L^2}$$

$$res \ _{z=+1} \ \frac{Q_0(z)}{z} = -\sqrt{(\frac{\epsilon\rho}{c} + \frac{\alpha}{c})^2 - m^2\rho^2c^2} \ ,$$

$$res \ _{z=-1} \ \frac{Q_0(z)}{z} = +\sqrt{(-\frac{\epsilon\rho}{c} + \frac{\alpha}{c})^2 - m^2\rho^2c^2}$$

$$res \ _{z=\infty} \frac{Q_0(z)}{z} = -res \ _{y=0} \frac{Q_0(y^{-1})}{y} = +\sqrt{\frac{\alpha^2}{c^2} - L^2}$$

1-term quantization provides us with

$$(\frac{1}{2}) \ (-2\pi i) \ \Bigg[2\sqrt{\frac{\alpha^2}{c^2} - L^2} + \sqrt{(\frac{\epsilon^2\rho^2}{c^2} - m^2\rho^2c^2 + \frac{\alpha^2}{c^2}) + 2\frac{\alpha\epsilon\rho}{c^2}} \ -$$

$$- \sqrt{(\frac{\epsilon^2\rho^2}{c^2} - m^2\rho^2c^2 + \frac{\alpha^2}{c^2}) - 2\frac{\alpha\epsilon\rho}{c^2}}\ \Bigg] = 2\pi\hbar\ n \tag{7.7b}$$

As for the next term contribution

$$Q_1 = -\frac{1}{2Q_0}\ [\ \frac{d}{dt}\ Q_0(t) + V(t)\ Q_0(t)\ ]\ . \tag{7.8a}$$

with (in units $c = 1, \hbar = 1$)

$$Q_0'(t) = \frac{+2e^{2t}}{(1-e^{2t})^2}\ \sqrt{\cdots}\ + + \ \frac{1}{(1-e^{2t})}\ \frac{1}{\sqrt{\cdots}}\times$$

$$\times \left[[2\epsilon e^t + \alpha(1+e^{2t})]\ (2\epsilon e^t + \alpha 2e^{2t}) - m^2 4e^{2t}\ +\ 2L^2(1-e^{2t})e^{2t} \right]\ ; \tag{7.8b}$$

for (Q_0'/Q_0) anf $V(t)$ one has

$$\frac{Q_0'}{Q_0} = \frac{2e^{2t}}{1-e^{2t}} + \frac{[2\epsilon e^t + \alpha(1+e^{2t})]\ (2\epsilon e^t + \alpha 2e^{2t})\ -\ m^2 4e^{2t}\ +\ 2L^2(1-e^{2t})e^{2t}}{[2\epsilon e^t + \alpha(1+e^{2t})]^2\ -\ m^2 4e^{2t}\ -\ L^2(1-e^{2t})^2}\ , \tag{7.8c}$$

$$V = \frac{1-e^{2t}}{e^{2t} + 2ae^t + 1} - \frac{1+e^{2t}}{1-e^{2t}}\ . \tag{6.8d}$$

Allowing for

$$\oint_{\mathcal{L}} Q_1(t)\ dt = -2\pi i\ \times$$

$$\times(-\frac{1}{2})\left\{ \sum_{z=0,\infty,\pm 1} res\ \left(\frac{1}{z}\frac{Q_0'(z)}{Q_0(z)}\right) + \sum_{z=0,\infty,\pm 1} res\ \left(\frac{1}{z}V(z)\right)\right\}\ . \tag{7.9}$$

we arrive at the vanishing contribution of the term Q_1. Therefore, 2-term quantization leads to

$$\epsilon = mc^2\ \left(1 + \frac{\alpha^2/\hbar^2c^2}{N^2}\right)^{-1/2}\ \sqrt{1 - \frac{\hbar^2}{m^2\rho^2c^2}\ (N^2 + \alpha^2/\hbar^2c^2)}\ . \tag{7.10}$$

where $N = n + \sqrt{k^2 - \alpha^2/\hbar^2c^2}$.

Consideration of the hydrogen atom on the base of Dirac equation in the space S_3 can be done in the same line. The final result is as follows

$$\epsilon = mc^2\ \left(1 + \frac{\alpha^2/\hbar^2c^2}{N^2}\right)^{-1/2}\ \sqrt{1 + \frac{\hbar^2}{m^2\rho^2c^2}\ (N^2 + \alpha^2/\hbar^2c^2)}\ . \tag{7.11}$$

where $N = n + \sqrt{k^2 - \alpha^2/\hbar^2c^2}$.

However, it must be stressed that these two formulas obtained are hardly correct in full because the second one (7.11) does not result in an exact result at $\alpha = 0$ (see Section 6) [3]. Moreover, turning to the next terms Q_2, Q_3 and so on , we might see that they give not-vanishing contributions. The latter indirectly proves only approximate character of the formula (7.1). In should be added that till now any exact wave function solutions of the latter task have not found.

---

[3] This formula was found in [68] on the base of other considerations, where it was considered as exact one.

References

1. Planck M. *Die Quantenhypothese für Molekeln mit mehreren Freiheitsgraden.* Verhandlungen der Deutschen Physikalischen Gesellschaft. 1915. Bd. 17. S. 407-418; S. 438-451.

2. Planck M. *Die physikalische Struktur des Phasenraum.* Annalen der Physik. 1916. Bd. 50. S. 385-418.

3. Wilson W. *The quantum theory of radiation and line spectra.* Phil. Magazine. 1915. Vol. 29. P. 795-802.

4. Ishiwara J. *Die universelle Bedeutung des Wirkungsquantums.* Tokio Sugaku Buturigakkawi Kizi. 1915. Bd 8. S. 106-116.

5. Sommerfeld A. *Zur Theorie der Balmerschen Serie.* Münchener Berichte. 1915. S. 425-458.

6. Sommerfeld A. *Die Feinstructur der wasserstoff und wasserstoffähnlichen Linien.* Münchener Berichte. 1915. S. 459-500; *Zur Quantentheorie der Spektrallinien.* Annalen der Physik. 1916. Bd 51. S. 1-94; S. 125-167.

7. Schwarzschild K. *Zur Quantenhypothese.* Berliner Berichte. 1916. S. 548-568.

8. Epstein P.S. *Zur Theorie des Starkeffektes.* Z. Phys. 1916. Bd. 17. S. 148-150.

9. Epstein P.S. *Zur Theorie des Starkeffektes.* Annalen der Physik. 1916. Bd. 50. S.489-520.

10. Epstein P.S. *Zur Quantentheorie.* Annalen der Physik. 1916. Bd. 51. S.168-188.

11. Sommerfeld A. *Atombau und Spektrallinien.* Braunchweig, Vieweg, 1919. S. 327-357; S. 520-522;

12. Wilson W. *The quantum theory and electromagnetic phenomena.* Proc. Roy. Soc. London. A. 1922. Vol. 102. P. 478-483.

13. Wentzel G. *Eine Verallgemeinerung der Quantembedingungen für die Zwecke der Wellenmechanik.* Z. Phys. 1926. Bd. 38. S. 518-529.

14. Kramers H.A. Z. Phys. 1926. Bd. 39. S. 828.

15. Brillouin M.L. *La méchanique ondulatoire de Schrödinger; une méthode générale de résolution par approximations successives.* Compt. Rend. Acad. Sci. Paris. 1926 Tom 183. P. 24-44.

16. Brillouin M.L. *Remarques sur la Méchanique ondulatoire.* J. Phys. Radium. 1926. Vol. 7. P. 353-368.

17. Dunham J.L. *The Wentzel-Brillouin-Kramers metod of solving the wave equation.* Phys. Rev. 1932. Vol. 41. P. 713-720.

18. Langer R.E. Bull. Amer. Math. Soc. 1934. Vol. 40. P. 545.

19. Langer R.E. *On the connection formulas and the solutions of the wave equation.* Phys. Rev. 1937. Vol. 51. P. 669-676.

20. Titchmarsh E.C. Quart. J. Math. 1954. Vol. 5. P. 228.

21. Leller J.B. *Corrected Borh-Sommerfeld quantum conditions for nonseparable systems.* Ann. Phys.

(N.Y.) 1958, Vol. 4, P. 180-188.

22. Bailey Paul B. *Exact quantization rules for the one-dimensional Scrödinger equation with turning points.* J. Math. Phys. 1964. Vol. 5, No 9. P. 1293-1297.

23. Froman N., Froman P.O. *JWKB Approximation: contributions to the theory.* North-Holland Publ. Co., Amsterdam, 1965.

24. Joseph B. Krieger. *Asymptotic properties of perturbation theory.* J. Math. Phys. 1966. Vol. 9, No 3. P. 432-435.

25. Ponomarev L.I. *Lectures in quasi-classics* (in Russian). Kiev. 1966.

26. Krieger J.B. Rosenzweig C. Phys. Rev. 1967. Vol. 164. P. 171.

27. Krieger J.B., Lewis M.L., Rosenzweig C. J. Chem. Phys. 1967. Vol. 47. P. 2942.

28. Krieger J.B. J. Math. Phys. 1968. Vol. 9. P. 432.

29. Rosenzweig C., Krieger J.B. *Exact quantization conditions.* J. Math. Phys. 1968. Vol. 9, No 6. P. 849-860.

30. Voros A. *Semi-classical approximations.* Ann. Inst. H. Poincaré. 1976. Vol. A24. P.31-90.

31. De Witt-Morette C. *The semiclassical expansion.* Ann. Phys.(N.Y.) 1976. Vol. 97. P. 367-399.

32. Neveu A. *Semiclassical quantization methods in field theory.* Phys. Rep. C. 1976. Vol. 23. P. 265-272.

33. Mizrahi M.M. *On the semiclassical expansion in quantum mechanics for arbitrary Hamiltonians.* J. Math. Phys. 1977. Vol. 18. P. 786-790.

34. Karl M. Bender, Kaare Olaussen and Paul S. Wang *Numerological analysis of the WKB approximation in large order.* Phys. Rev. D. 1977. Vol. 16, No 6. P. 1740-1748.

35. Gallas J.A.C. Chem. Phys. Lett. 1983. Vol. 96, No 1. P. 479-480.

36. Comtet A., Bandrauk A.D., Campbell D.K. *Exactness of semiclassical bound state energies for supersymmetric quantum mechanics.* Phys. Lett. B. 1985. Vol. 150. P. 159-162.

37. Dutt R., Khare A., Sukhatme U.P. *Exactness of supersymmetric WKB spectra for shape-invariante potentials.* Phys. Lett. B. 1986. Vol. 181. P. 295-298.

38. Gomes M.A.F., Thomaz M.T., Vasconcelos G.L. *Matrix formulation for the Wentzel-Kramers-Brillouin quantization rule.* Phys. Rev. A. 1986. Vol . 34, No 5. P. 3598-3604.

39. Schöpf Hans-Georg. *Zur Geschichte der Bohr-Sommerfeldschen Quantentheorie.* Ann. Phys. 1988. Bd. 45, No 8. S. 595-604.

40. Kobylinsky N.A., Stepanov S.S., Tutik R.S. *Semiclassical approach to ground state within the Klein-Gordon equation.* J. Phys. A. 1990. Vol. 23, No 6. P. 237-241.

41. Barclay D.T., Khare A. Sukhatme U. *Is the lowest order supersymmetric WKB approximation exact for all shape invariant potentials?* – 8 pages. hep-th/9311011.

42. Funahashi K., Kashiwa T., Sakoda S., Fujii K. *Exactness in the Wentzel-Kramers-Brillouin approximation for some homogeneous spaces.* J. Math. Phys. 1995. Vol. 36. P. 4590-4611; hep-th/9501145.

43. De Vega H.J., Larsen A.L., Sanchez N. *Semi-classical quantization of circular strings in de Sitter and anti de Sitter spacetimes.* Phys. Rev. D. 1995. Vol. 51. P. 6917-6928; hep-th/9410219.

44. Robnik M., Salasnich L. *WKB to all orders and the accuracy of the semiclassical quantization.* J. Phys. A. 1997. Vol. 30. P. 1711-1718; quant-ph/9610027.

45. Robnik M., Salasnich L. *WKB exactness for the angular momentum and the Kepler problem: from the torus quantization to the exact one.* J. Phys. A. 1997. Vol. 30. P. 1719-1729; quant-ph/9603014.

46. Schrödinger E. *A method of determining quantum-mechanical eigenvalues and eigenfunctions.* Proc. Roy. Irish. Soc. A. 1940. Vol. 46, No 1. P. 9-16.

47. Infeld L., Schild A. *A note on the Kepler problem in a space of constant negative curvature.* Phys. Rev. 1945. Vol. 67, No 2. P. 121-122.

48. Bander M., Itzykson C. *Group theory and the hydrogen atom. I,II* Rev. Mod. Phys. 1966. Vol. 38. P. 330-345; P. 346-357.

49. Aronson E.B., Malkin I.A., Manko V.I. *Dynamic symmetries in quantum theory.* (in Russian). Elementary Particles and Atomic Nuclei. 1974. Vol. 5, No 1. P. 122-171.

50. Higgs P.W. *Dynamical symmetries in a spherical geometry. I.* J. Phys. A. 1979. Vol. 12, No 3. P. 309-323.

51. Leemon H.I. *Dynamical symmetries in a spherical geometry. II.* J. Phys. A. 1979. Vol. 12, No 14. P. 489-501.

52. Ringwood G.A. Devreese J.T. *The hydrogen atom: Quantum mechanics on the quotient of a conformally flat manifold.* J. Math. Phys. 1980. Vol 21. P. 1390-1392.

53. Kurochkin Yu.A., Otchik V.S, *Analogue of the Gunge-Lenz vector and energy spectrum in Kepler problem on 3-sphere.* Doklady AN BSSR. 1979. Vol. 23, No 11. P. 987-990.

54. Bogush A.A., Kurochlin Yu.A., Otchik V.S. *On quantum-mechanical Kepler problem in Lobachevsky space.* Doklady AN BSSR. 1980. Vol. 24, No 1. P. 19-22.

55. Bessis N., Bessis G., Shamseddine R. *Atomic fine structure in a space of constant curvature.* J. Phys. A. 1982. Vol. 15, No 10. P. 3131-3144.

56. Xu Chondming , Xu Dianyan. *Dirac equation and energy levels of hydrogen-like atoms in Robertson-Walker metric.* Nuovo Cim. B. – 1984. Vol. 83, No 2. P. 162-172.

57. Bessis N., Bessis G., Shamseddine R. *Space-curvature effects in atomic fine- and hyperfine-structure calculations.* Phys. Rev. A. 1984. Vol. 29, No 5. P. 2375-2383.

58. Duval C., Künzle H. *Minimal gravitational coupling in the Newtonian theory and the covariant Schrödinger equation.* Gen. Rel. Grav. 1984. Vol. 16. P. 333-347.

59. Shamseddine R., *Structure fine et hyperfine atomique dans un espace à courbure constante.* J. Phys. A. 1986. Vol. 19, No 5. P. 717-724.

60. Bessis N., Bessis G., Roux D. *Space-curvature effects in the interaction between atom and external fields: Zeeman and Stark effects in a space of constant positive curvature.* Phys. Rev. A. 1988. Vol. 33, No 1. P. 324-336.

61. Tagirov E.A. *Quantum mechanics in Riemannian space-time. General covariant Schrödinger equation with relativistic corrections.* Dubna, 1988. Preprint / E2-88-678.

62. Gorbatsievich A.K., Priebe A. *On the hydrogen atom in Kerr space time.* Acta Phys. Polon. B. 1989. Vol. 20, No 11. P. 901-909.

63. Groshe C. *The path integral for the Kepler problem on the pseudosphere.* Ann. Phys. (N.Y.) 1990. Vol. 204. P. 208-222.

64. Katayama N. *A note on the Kepler problem in a space of constant curvature.* Nuovo Cim. B. 1990. Vol. 105, Ser. 2, No 1. P. 113-119.

65. Foerster A., Girotti H.O., Kuhn P.S. *Nonrelativistic quantum particle in a curved space as a constrained system.* – hep-th/9411052.

66. Gorbatcevich A.K., Tomilchik L.M. *Equation of motion of particles in conformally-flat space and quark confinement.* Minsk, 1986. 9 pages. Preprint No 415, Belarus Academy of Siences, Institute of physics.

67. Melnikov V.N. Shikin G.N. *Hydrogen-like atom in gravitational field of the universe.* Izvestiya Vuzov. Fizika. 1985. No 1. P. 55-59.

68. Shamseddine R. *On the resolution of the wave equations of electron in a space of constant curvature.* Can. J. Phys. 1997. Vol. 75, No 11. P. 805-811.